

Magnitude of HIV Index Case Testing among HIV Positive Clients Attending ART Clinic in University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia, 2023

Lakew Asmare^{1*}, Zeamanuel Yigzaw², Tihtna Alemu³

¹Department of Epidemiology and Biostatistics, School of Public Health, College of Medicine and Health Sciences, Wollo University, Dessie, Ethiopia

²Department of Health Promotion and Behavioral Science, School of Public Health, College of Medicine and Health Sciences, Bahir Dar University, Bahir Dar, Ethiopia

³Department of Surgical Nursing, School of Nursing, College of Medicine and Health Sciences, University of Gondar, Ethiopia

*Correspondence should be addressed to Lakew Asmare, lakewt071981@gmail.com

Received date: January 25, 2024, **Accepted date:** March 07, 2024

Citation: Asmare L, Yigzaw Z, Alemu T. Magnitude of HIV Index Case Testing among HIV Positive Clients Attending ART Clinic in University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia, 2023. J AIDS HIV Treat. 2024;6(1):37-43.

Copyright: © 2024 Asmare L, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Index case testing is the provision of HIV testing services to family members or sexual partners of individuals already diagnosed with HIV. Its promotion stems from its potential to detect new cases, improve testing coverage, enhance ART retention, and provide convenience to affected families. Although evidence on index case testing in Ethiopia, particularly in the study area, is lacking, even, it is regarded as a crucial strategy for reducing HIV transmission among those unaware of their HIV status.

Objective: This study aimed to determine the prevalence and associated factors with HIV index testing in Northwest Ethiopia.

Methods: The study utilized an institution-based cross-sectional study design. The recruitment of 387 HIV-positive clients was done using a simple random sampling technique, and structured interviewer-administered questionnaires were employed for data collection. The collected data was entered into Kobo Collect software and later exported to STATA version 16 for analysis. In binary logistic regression, variables with a p-value below 0.25 were considered as candidates for multivariate logistic regression. An association was assessed if the p-value was below 0.05 with a 95% confidence interval.

Results: In the study, 380 (98.19% of participants) underwent the interview. The prevalence of HIV index case testing service was 38.7% (CI: 36.45% to 42.03%). Significant associations were observed between HIV index case testing and certain factors. Those with a higher education level (college and above) had 1.21 times higher odds (AOR) (CI: 1.05 to 3.11). Married individuals had 2.30 times higher odds of receiving index case testing (AOR) (CI: 1.98 to 4.20). Participants engaged in work for monetary compensation had 1.66 times higher odds of undergoing index case testing (AOR) (CI: 1.58 to 4.71).

Conclusion: The utilization of index case testing service was lower in this study compared to previous research. Higher education level (college and above), occupation involving work for monetary compensation, and being married were associated with HIV index case testing. To enhance uptake, community-based education programs and targeted training for HIV-positive individuals are recommended. These measures can raise awareness and encourage greater participation in index case testing.

Keywords: HIV Counseling, HIV index case testing, HIV infection, HIV prevention, University of Gondar Specialized Hospital

Introduction

The Human Immunodeficiency Virus (HIV) and its sequelae, acquired immunodeficiency syndrome (AIDS), remain a significant global public health challenge. According to the World Health Organization [1], approximately 38.4 million people globally were living with HIV in 2021, with 1.5 million new infections and 650,000 AIDS-related deaths. Antiretroviral therapy (ART) has been instrumental in managing HIV, improving quality of life, and achieving viral suppression, thereby preventing onward transmission [2]. In 2020, there were an estimated 1.5 million people newly infected with HIV and 67.5% were in sub-Saharan Africa [3].

HIV testing and counseling was the gateway to ART, reducing morbidity and mortality and motivating individuals to change risky behaviours, as well as preventing infection by suppressing viral load in HIV-infected people [4,5]. To achieve the UNAIDS Project 95-95-95 objectives, which aim to ensure that 95% of people with HIV know their HIV status, 95% of those diagnosed with HIV are on antiretroviral therapy (ART), and 95% of those on ART achieve virologic control, alternative and cost-effective methods are needed. These methods include partner profiles and index partner-testing strategies, which can identify the HIV status of 95% of individuals with HIV. These strategies specifically target those who are on ART, ensuring that they receive the necessary support and interventions to achieve viral suppression and maintain good health [6-8]. In response to the devastating impact of HIV/AIDS, the government of Ethiopia has taken action to prevent further transmission and control the spread of the disease. The objective of the index case trial is to interrupt the chain of HIV transmission by promptly connecting HIV-negative individuals who are identified as being at risk to appropriate care and treatment in the event of HIV infection. This approach aims to effectively address the spread of HIV/AIDS and improve the overall health outcomes of affected individuals [9-11]. HIV index case testing is advocated for its ability to yield a high number of positive cases, enhance testing coverage, improve retention in antiretroviral therapy (ART), and provide a convenient service for families affected by HIV infection [10,12,13]. On a national scale, the proportion of males and females who underwent HIV testing among index cases in 2016 was 78.95% and 78.3%, respectively [14]. Demand generation and novel testing approaches will be necessary to reach undiagnosed people living with HIV and to promote frequent retesting among sexual contact of index case [11,15]. Because it has high yield positivity, improves testing coverage, improves ART retention, and offers a convenient service for families affected by HIV irrespective of the background of HIV prevalence in the country [16].

Despite the effectiveness of index case testing in identifying individuals at high risk of acquiring HIV and yielding positive cases, there is a significant lack of participation in Ethiopia, with many index clients failing to involve their partners and

biological children in testing. Therefore, the aim of this study was to assess the index case testing and its associated factors in the study area. And also used for empirical evidence of HIV/AIDS index case testing magnitude.

Methods

Study area and period

The study was conducted in University of Gondar Comprehensive Specialized Hospital, ART clinic from April 20/2023 to May 20/2023. Gondar Comprehensive Specialized Hospital is located in Gondar city, which is 735 km far from Addis Ababa, capital city of Ethiopia. There are ten governmental facilities in Gondar (eight health facilities and two hospitals) [17]. According to the hospital administration head report, University of Gondar Comprehensive Specialized Hospital serves nearly 250,000 patients in outpatient services; of this about 4,244 adult patients were ART users.

Study design

An Institution based cross-sectional study design was conducted.

Source population: All HIV positive adults who were attending ART clinic in Gondar Comprehensive Specialized Hospital.

Study population

All HIV positive adults >18 years who were receiving anti-retro viral therapy during data collection period.

Sample size determination

It was determined by using single population proportion formula by taking $p=50\%$, because there is no previously similar study done in the country, margin of error, $d=0.05$ and 95% confidence interval. So, the calculated sample size was 384. Since the source population is less than 10,000, $N=4,244$ using finite population correction formula we calculated the final sample size $n_f = \frac{n}{1+\frac{n}{N}}$, then, it gives the sample size of 387 adult HIV positive clients with 10% non-response rate.

Sampling procedure

Systematic random sampling technique was used by considering the average number of adult HIV positive clients visit in this hospital per month and the desired number of sample size to determine the interval of respondent was interviewed (k). Based on the registration book and ART focal health professional report, on average 25 clients visited this hospital per day. Therefore, with 20 working days in a month, 500 outpatients visit monthly and led to take every $k=2^{\text{nd}}$ interval of clients from ART clinic by selecting the first patient by lottery method.

Measurements

HIV index case testing: HIV testing for partner or family or any other sexual partner. Which was measured as “Yes” for did your HIV status elicit by index case contact? [18]

Index cases: Individuals diagnosed HIV positive that aware already enrolled in HIV care and treatment center [19].

Waiting time to HIV test result: The time taken to produce either positive or negative result [20].

Testing place: Index case testing service took place at community or facility [21].

Data collection procedure: Structured questionnaire was developed by referring different literatures from standard HIV index case testing tool [11,19,20,22-24]. The interview was translated into Amharic language then it was translated back to English to ensure consistency of questions. The questionnaire was separated into socio demographic, HIV index case testing, related variables, and other variables. A data collector was selected from health professionals who were trained on HIV testing program and with previous experience in data collection. Two diploma nurses were assigned as data collectors and one BSc nurse was assigned as a supervisor.

Data quality control: To ensure the quality of data, pretesting was done among 19 (5%) adult HIV positive clients at Debar

ART clinic. A two-day training was given about questionnaire. Prior to analysis, data was cleaned up and cross-checked. The principal investigator and the supervisor closely monitored the process throughout the data collection period and made due corrections.

Data management and analysis: The filled questionnaires were exported from open data kit/ODK to STATA version 16 for further analysis. Descriptive and summary statistics were presented in the form of text and tables. In binary logistic regression, variables with a p-value below 0.25 were considered as candidates for multivariate logistic regression. An association was assessed if the p-value was below 0.05 with a 95% confidence interval. Multi-collinearity was checked using variance inflation factor (VIF), which was 1.68. We assessed the goodness of fit for the corresponding model using the Pearson goodness-of-fit which was insignificant (p-value=0.16), therefore our model is a good fitted model.

Results

Among the total study participants, 380 (98.19%) were involved in the interview. Regarding their religion, 320 (84.21%) were Orthodox Christians, and 316 (83.83%) of them were urban residents. Concerning their educational status, the majority, 106 (27.89%) of them attended secondary education (**Table 1**).

Table 1. Socio-Demographic Characteristics of study participants in University of Gondar Comprehensive Specialized Hospital in Northwest Ethiopia in 2023 (n=380).

Variables	Category	Frequency	Percentage
Age	19-30	76	20
	31-40	105	27.63
	41-50	112	29.17
	≥ 51	87	22.84
Sex	Male	161	42.36
	Female	219	57.64
Place of residence	Urban	316	83.83
	Rural	64	16.17
Religion	Muslims	56	14.74
	Orthodox	320	84.21
	Others	4	1.05
Educational status	Unable to read & write	89	23.42
	Able to read & write	37	9.73
	Primary school	82	21.57
	Secondary school	106	27.89
	College & above	56	14.73
Marital status	Married	178	46.82
	Unmarried	202	53.18

Occupational status	Farmer	148	38.94
	Governmental	103	27.10
	Housewife	107	28.15
	Others	12	3.15
Work for money	No	71	18.69
	Yes	309	81.31
Sex with spousal partner	No	147	38.68
	Yes	233	61.32
Others=Merchant, Student, Private.			

HIV index case testing and other characteristics

This study assessed the magnitude of HIV index case testing, only 39.21% (CI=36.45%, 42.03%) of the respondents. Regarding waiting time to HIV test result 281 (73.95%) said 15-30 minute, and 305 (80.30%) of them were tested at the facility. The majority 324 (85.26%) of them were taking ART to manage HIV, and 242 (63.68%) were badly treated because they have HIV (**Table 2**).

Associated factors with index case testing

In this model, variables such as sex marital status, sex with spousal, education, work for money, as well as testing place were included in multivariable regression since their overall p-value in bi-variable regression was less than 0.25. On multiple logistic regression analysis, three variables were found to be associated with the HIV index case testing, which were work for money, education, and marital status.

Thus, our study showed participants who had educational status, college & above increased the odds of HIV index testing

by a factor of 1.21 times (AOR=1.21, 95%, CI, 1.05, 3.11) as compared to those who can't read and write. Another finding also suggested that those who were married increased the odds of HIV index testing by 2.30 times (AOR=2.30, 95%, CI, 1.98, 4.20) as compared to their counterpart. Additionally, study participants who work for money increased 1.66 times HIV index testing (AOR=1.66, 95%, CI, 1.58, 4.71) as compared their counterpart (**Table 3**).

Discussion

This study assessed HIV index case testing service in the University of Gondar hospital. The results showed that only 39.21% (CI=36.45%, 42.03%) of the respondents got the HIV index case. This means that HIV index case testing service provided to the index case client was in a substandard way. As a result, the transmission of HIV becomes high.

This study is inline with study done in Lesotho, which was 37.3% [24]. The possible justification might be having similar infrastructure and health system policy regarding HIV testing service but, the finding is lower than the WHO standard,

Table 2. HIV index case testing, related variables, and other variables in the University of Gondar Comprehensive Specialized Hospital in Northwest Ethiopia, 2023 (n=380).

Variables	Category	Frequency	Percentage
Waiting time to HIV test result	15-30 minute	281	73.95
	31-60 minute	61	16.05
	61-90 minute	26	6.84
	91-max minute	12	3.16
Testing place	Community	75	19.70
	Facility	305	80.30
Taking ART to manage HIV	Yes	324	15
	No	56	85
heard about badly treated	Yes	236	63
	No	144	37
Index case test their contact	No	231	61.49
	Yes	149	39.21

Table 3. Multiple logistic regression of HIV index case testing in in University of Gondar Comprehensive Specialized Hospital in Northwest Ethiopia.in 2023(n=380).

Variables	Categories	Index case testing		COR (95%CI)	AOR (95%CI)
		Yes	No		
Sex	Male	80	81	0.46 (0.04, 0.950)	0.15 (0.07, 2.4)
	Female	69	150	1.00	1.00
Mstatus	Married	41	137	3.83 (1.2, 9.14) **	2.30 (1.98, 4.20) *
	Unmarried	108	94	1.00	1.00
Spousal sex	Yes	81	152	1.61 (0.91, 5.04)	1.40 (0.97, 3.91)
	No	68	79	1.00	1.00
Education	College & above	26	30	0.35 (0.75, 2.40) **	0.21 (0.15, 0.91) *
	Secondary	44	62	0.43 (0.75, 2.40)	0.15 (0.05, 3.01)
	Primary	28	54	0.59 (0.51, 1.53)	0.45 (0.75, 2.28)
	Able to read &write	30	17	0.17 (1.15, 2.42)	0.21 (1.17, 52)
	Unable to read &write	21	68	1.00	1.00
Work for money	Yes	99	210	5.05 (1.27, 8.36) **	1.66 (1.58, 4.71) *
	No	50	21	1.00	1.00
Testing place	Community	50	25	0.24 (0.11, 1.67)	0.21 (0.01, 0.67)
	Facility	99	206	1.00	1.00

Note: COR: Crude Odd Ratio; AOR: Adjusted Odd Ratio

the National HIV index case testing strategy of Ethiopia and PEPFAR guideline [24,25]. The possible explanations might be due to HIV index case testing in the current study could be the nature of the HIV intervention, as it is a complex intervention and resource-intensive demanding trained providers. Hence, resource limitation, lack of training of the ICT providers, budget constraints, and lack of NGOs support in the study area could be the potential reasons. This finding is also relatively similar from a study conducted in Eswatini, where index case did not got the index case testing service according to standard [25]. A possible reason for this similarity could be the similarity of the health system infrastructure, the presence of the same methodology and dimensions used to measure index case testing.

Thus, our study showed participants who had educational status, college & above increase the odds of HIV index testing by a factor of 1.21 times (AOR=1.21, 95% CI, 1.05, 3.11) as compared to those who can't read and write. This finding were supported by a study done in Gamo Gofa and study done in SNNPR [24,26]. This is because, when educational status increases, client responsiveness on index case testing service also increases this may be due to those who are attending higher education may have a variety of experiences. Another finding also suggested that those who were married increases the odds of HIV index testing by 2.30 times (AOR=2.30, 95% CI, 1.98, 4.20) as compared to their counterparts. This finding is supported by a study done in USA, where married

individuals were 4.3 times more likely involve in HIV index testing service than unmarried individuals [27] and a study done in western Kenya, where single/never married persons were 13 times as less likely to involve in HIV/AIDS service than married individuals [28]. This might be due to the fact that, HIV clients who are married disclose their HIV status to others and they have a tendency to notify their sexual clients/partners. Additionally, study participants who work for money increases 1.66 times HIV index testing (AOR=1.66, 95% CI, 1.58, 4.71) as compared to their counterparts. It might be low socio-economic status such as lack of transportation to health facility, poverty, and poor health, ultimately affect economic status of individual.

Strength and imitations of study

This study was conducted in a less studied population. It is the first study of its kind in the study area with a validated and comprehensive tool, which provides updated information. And also, it may serve as the basis for future studies and may offer suggestions for strengthening HIV index case testing service that can be used as an input for improvement of HIV transmission.

However, self-reported information is subjected to reporting errors, or there might be social desirability bias, and this may lead to overestimate/underestimate the finding.

Conclusion

Factors such as having a college education or higher, engaging in work for monetary compensation, and being married were associated with higher rates of HIV index case testing. To improve the uptake of HIV index case testing, it is recommended to implement community-based education programs, provide training for healthcare professionals, and raise awareness among individuals receiving antiretroviral therapy (ART). These measures can contribute to enhancing HIV index case testing rates.

Abbreviations

ART: Anti-Retroviral Therapy, AOR: Adjusted odd ratio, ARV: Anti-Retroviral, CDC: Centers of Disease Control and Prevention, CI: Confidence Interval, COR: Crude odd ratio, FSWs: Female Sex Workers, HIV: Human immune-deficiency virus, HTS: HIV Testing Service, ICT: Index Case Testing

Declaration

Ethical approval was obtained from the institution review board of the University of Gondar, Institute of Public Health. Oral informed consent was obtained from participants in the Ethical approval process. Confidentiality of information and privacy of participants were assured for all the information provided. All the collected information was put to assure personal confidentiality using digital data collection technique and locked by password for digital instruments. Informed consent was completely volunteer, and participants could withdraw or refuse at any time during the process and their data was used only for research purposes. In case the participants were unable to read and write, the guardians were involved during informed consent.

Consent for Publication

Not applicable.

Availability of Data and Materials

All necessary data are included in this manuscript.

Competing Interest

Authors declare that they have no conflict of interest.

Funding

No funding was obtained for this study.

Authors' Contribution

Conceptualization: LA & ZY, data curation: LA & TA, formal analysis: LA & TA, investigation: LA & CA, methodology: LA, software: LA, validation: LA & ZY, visualization: LA, ZY, & TA,

writing original draft: LA, writing review & editing: LA, ZY, &TA. All authors contributed to the article and approved the submitted version accordingly based on the above descriptions.

Acknowledgements

We would like to thank the University of Gondar and Institute of public health, Department of health policy and System as well as the University of Gondar Comprehensive Specialized Hospital for their cooperation and permission to conduct the study. We are also grateful to all the respondents, data collectors, and supervisors for providing valuable time and necessary information for this research.

References

1. Arantes LMN, Pedrosa AO, Meneguetti MG, Gir E, Botelho EP, Silva ACdOe, et al. Factors Associated with Late Diagnosis of Human Immunodeficiency Virus/Acquired Immunodeficiency Syndrome (HIV/AIDS) in a University Hospital in Brazil: Challenges to Achieving the 2030 Target. *Viruses.* 2023;15(10):2097.
2. Koduri R. Factors Associated with Quality of Life Among People Living with HIV In India: University of Pittsburgh; 2023.
3. Little K, Thorne C, Luo C, Bunders M, Ngongo N, McDermott P, et al. Disease progression in children with vertically-acquired HIV infection in sub-Saharan Africa: reviewing the need for HIV treatment. *Current HIV Research.* 2007;5(2):139-53.
4. Robbins RN, Spector AY, Mellins CA, Remien RH. Optimizing ART adherence: update for HIV treatment and prevention. *Current HIV/AIDS Reports.* 2014;11:423-33.
5. Granich R, Gupta S, B Suthar A, Smyth C, Hoos D, Vitoria M, et al. Antiretroviral therapy in prevention of HIV and TB: update on current research efforts. *Current HIV research.* 2011;9(6):446-69.
6. Amstutz A. Supporting Lesotho on the way towards the UNAIDS 90-90-90 targets: Operational and clinical research addressing HIV/AIDS care in resource-limited settings. University of Basel Associated Institution; 2021.
7. Barker JL. Effective Public Health Strategies to Optimize HIV Testing Services Among Men Ages 15 Years and Older in Kenya: The University of North Carolina at Chapel Hill; 2023.
8. Okoboi S. Peer distribution of HIV self-test kits to men who have sex with men in Uganda: University of Antwerp; 2021.
9. Organization WH. Policy brief: Consolidated guidelines on HIV prevention, diagnosis, treatment and care for key populations. World Health Organization; 2017.
10. Dougherty G, Abena T, Abesselo JP, Banda JN, Biyaga TP, Boccanera R, et al. Improving services for HIV-exposed infants in Zambia and Cameroon using a quality improvement collaborative approach. *Global Health: Science and Practice.* 2021;9(2):399-411.
11. WHO. Guidelines on HIV self-testing and partner notification: supplement to consolidated guidelines on HIV testing services:

World Health Organization; 2016.

12. Patel HK, Duong YT, Birhanu S, Dobbs T, Lupoli K, Moore C, et al. A comprehensive approach to assuring quality of laboratory testing in HIV surveys: lessons learned from the population-based HIV impact assessment project. *J AIDS Journal of Acquired Immune Deficiency Syndromes.* 2021;87:S17-S27.

13. Saito S, Howard AA, Chege D, Ellman TM, Ahoua L, Elul B, et al. Monitoring quality at scale: implementing quality assurance in a diverse, multicountry HIV program. *AIDS (London, England).* 2015;29(Suppl 2):S129.

14. McDaid D, Wiley M, Maresso A, Mossialos E, Organization WH. Ireland: Health system review. 2009.

15. Bekele YA, Fekadu GA. Factors associated with HIV testing among young females; further analysis of the 2016 Ethiopian demographic and health survey data. *PLoS one.* 2020;15(2):e0228783.

16. Mirzajani H, Mahmud R, Fauzi Mohd Ayub A, Wong SL. Teachers' acceptance of ICT and its integration in the classroom. *Quality Assurance in Education.* 2016;24(1):26-40.

17. Zemariam AB, Anlay DZ, Alamaw AW, Abebe GK, Techane MA. Incidence and Predictors of Major Adverse Drug Reactions Among Human Immunodeficiency Virus-Infected Children on Antiretroviral Treatment in West Amhara Comprehensive Specialized Hospitals, Northwest Ethiopia: A Multicenter Retrospective Follow-up Study. *Clinical Therapeutics.* 2023.

18. Berlinger N, Wynia M, Powell T, Hester DM, Milliken A, Fabi R, et al. Ethical framework for health care institutions responding to novel Coronavirus SARS-CoV-2 (COVID-19) guidelines for institutional ethics services responding to COVID-19. *The Hastings Center.* 2020;12(3):1-12.

19. Edosa M, Merdassa E, Turi E. Acceptance of Index Case HIV Testing and Its Associated Factors Among HIV/AIDS Clients on ART Follow-Up in West Ethiopia: A Multi-Centered Facility-Based Cross-Sectional Study. *HIV/AIDS-Research and Palliative Care.* 2022;451-60.

20. Greenwald JL, Burstein GR, Pincus J, Branson B. A rapid review of rapid HIV antibody tests. *Current infectious disease reports.* 2006;8(2):125-31.

21. WHO. Guidelines: updated recommendations on HIV prevention, infant diagnosis, antiretroviral initiation and monitoring. 2021.

22. Williams D, MacKellar D, Dlamini M, Byrd J, Dube L, Mndzebele P, et al. HIV testing and ART initiation among partners, family members, and high-risk associates of index clients participating in the CommLink linkage case management program, Eswatini, 2016–2018. *Plos one.* 2021 Dec 20;16(12):e0261605.

23. Endalamaw A, Geremew D, Alemu SM, Ambachew S, Tesera H, Habtewold TD. HIV test coverage among pregnant women in Ethiopia: A systematic review and meta-analysis. *African Journal of AIDS Research.* 2021;20(4):259-69.

24. Jubilee M, Park FJ, Chipango K, Pule K, Machinda A, Taruberekera N. HIV index testing to improve HIV positivity rate and linkage to care and treatment of sexual partners, adolescents and children of PLHIV in Lesotho. *PLoS One.* 2019;14(3):e0212762.

25. Armstrong-Mensah EA, Tetteh AK, Ofori E, Ekhosuehi O. Voluntary counseling and testing, antiretroviral therapy access, and HIV-related stigma: global progress and challenges. *International Journal of Environmental Research and Public Health.* 2022;19(11):6597.

26. Luo M, Hann K, Zhang G, Pan X, Jiang J, Chen L, et al. HIV testing uptake and yield among sexual contacts of HIV-positive men who have sex with men in Zhejiang Province, China, 2014–2016: a cross-sectional pilot study of a choice-based partner tracing and testing package. *PLoS One.* 2020 Jun 4;15(6):e0232268.

27. Kposowa AJ. Marital status and HIV/AIDS mortality: evidence from the US National Longitudinal Mortality Study. *International Journal of Infectious Diseases.* 2013;17(10):e868-e74.

28. Sharma M, Kariithi E, Kemunto E, Otieno G, Lagat H, Wamuti B, et al. High acceptability of assisted partner notification services among HIV-positive females in Kenya: results from an ongoing implementation study. *Journal of Acquired Immune Deficiency Syndromes.* 2021;86(1):56.